

## Minor Skin Surgery for Skin Lesions Commissioning Policy

Interventions	<b>Minor Surgery for Skin Lesions</b>
Policy Criteria	<p>Skin Lesions must meet at least ONE of the following criteria to be removed<sup>3</sup>:</p> <ul style="list-style-type: none"> <li>• The lesion is unavoidably and significantly traumatised on a regular basis (e.g. causing regular bleeding or recurrent infections). There is repeat infection requiring 2 or more antibiotics per year</li> <li>• The lesion bleeds in the course of normal everyday activity</li> <li>• The lesion is obstructing an orifice or impairing visual access. The lesion significantly impacts on function eg: restricts joint movement</li> <li>• If left untreated, more invasive intervention would be required for removal</li> <li>• Facial viral warts that have not resolved with an appropriate trial of topical treatment.</li> <li>• Facial spider naevi in children causing significant psychological impact</li> </ul>
Background	<p>NHS Scarborough and Ryedale CCG and NHS Vale of York CCG are responsible for commissioning activity in secondary care, and this policy sets out the criteria for referral to secondary care for minor surgery, as this is not always routinely commissioned.</p> <p>As well as the lesions specifically detailed in the policy, the policy also applies to the benign lesions listed below<sup>3</sup>:</p> <p>Please note: This list is not exclusive:</p> <ul style="list-style-type: none"> <li>• Solar comedones</li> <li>• Corn/callous</li> <li>• Dermatofibroma</li> <li>• Milia</li> <li>• Epidermoid &amp; Pilar Cysts (sometimes incorrectly called sebaceous cysts)</li> <li>• Seborrheic keratosis (basal cell papillomata)</li> <li>• Spider naevi (telangiectasia)</li> <li>• Xanthelasmata</li> <li>• Neurofibromata</li> </ul>

<p>Commissioning Position</p>	<p><b>Treatment of any condition for purely cosmetic reasons is not commissioned.</b></p> <p>NHS Scarborough and Ryedale CCG and NHS Vale of York CCG only commission referrals to secondary care dermatology / plastic surgery in the following circumstances:</p> <ul style="list-style-type: none"> <li>• Where there is diagnostic uncertainty or a possibility of malignancy</li> </ul> <p>OR</p> <ul style="list-style-type: none"> <li>• A lesion has been excised in primary care and a re-excision has been subsequently recommended on clinical grounds by the histopathologist</li> </ul> <p>OR</p> <ul style="list-style-type: none"> <li>• After individual approval by the Individual Funding Request Panel (IFR)</li> </ul> <p>The following conditions should always be referred direct to secondary care (dermatology, head and neck surgery or plastic surgery as appropriate) and IFR approval is not required for:</p> <ul style="list-style-type: none"> <li>• Malignant Melanoma (2 week pathway)</li> <li>• Squamous Cell Carcinoma (SCC) including extensive premalignant changes to the lip (2 week pathway)</li> <li>• Basal Cell Carcinoma (refer as <b>urgent and not via 2 week pathway. Where possible those &lt;1cm and below the clavicle should be excised in Primary Care</b>).</li> </ul> <ul style="list-style-type: none"> <li>○ Removal by accredited GP Minor Surgeon (either in-house or through Practice-to-Practice referral via LES scheme)</li> <li>○ Remove with 4mm margins, send for histology</li> </ul> <ul style="list-style-type: none"> <li>• Lentigo Maligna</li> <li>• Naevus Sebaceous</li> </ul>
<p>Indications</p>	<p><b>Criteria for secondary care referral</b></p>
<p>Benign Skin Lesions</p>	<p>The removal of benign skin lesions is not routinely commissioned for cosmetic reasons.</p> <p><b>Where there is diagnostic uncertainty GPs should send three photos, (field, close-up and dermatoscopic) to the Dermatologists for advice on whether the patient needs to be seen in secondary care or whether primary care excision biopsy is appropriate (“permission to biopsy”)</b></p> <p>Under the Minor Surgery Directed Enhanced Service, GP practices may undertake:</p>

	<ul style="list-style-type: none"> <li>• Incision and drainage of an abscess requiring local anaesthetic</li> <li>• Excision of sebaceous cysts where there is a history of more than one infection</li> <li>• Incision and Curettage of Meibomian Cysts (as per the Commissioning Statement <a href="#">Click Here</a>)</li> </ul> <p><b>Referral to Secondary Care services</b></p> <p>Indications for referral to an appropriate alternative provider include:</p> <ul style="list-style-type: none"> <li>• lesions suspicious of being a basal cell carcinoma (BCC) that are &gt; 1cm in size or above the clavicle or squamous cell carcinoma (SCC) and melanomas.</li> <li>• lesions of uncertain significance where a specialist opinion is that primary care treatment is appropriate or a histological diagnosis is required that should be seen and managed by an accredited clinician who has links with the local skin cancer MDT. This would include secondary care dermatologists and also (where commissioned) GPwSIs.</li> <li>• sebaceous cysts where there is a history of one or more episodes of infection and so which would be appropriate for removal under this enhanced service, but where the <ul style="list-style-type: none"> <li>○ patient has a history of keloid scarring or hypertrophic scarring and the lesion is in an area where the patient would not want to risk the development of such scarring</li> </ul> </li> </ul> <p><b>OR</b></p> <ul style="list-style-type: none"> <li>○ where the lesion lies in a position which is not appropriate for removal in primary care e.g. face or centre of spine</li> </ul> <p>All other requests must have prior approval through Individual Funding request Panel.</p>
Molluscum contagiosum	<p>Patients need to be managed in primary care. Referral to the dermatology department should only be made if patients have either of the following:</p> <ul style="list-style-type: none"> <li>• molluscum contagiosum in immunosuppressed patients</li> </ul>

	<ul style="list-style-type: none"> <li>• Diagnostic uncertainty of a solitary lesion.</li> </ul> <p>All other requests for referral for secondary care should have prior approval from individual funding request panel. Funding for treatment will not normally be commissioned.</p> <p>Where molluscum contagiosum is causing significant problems in the management of atopic eczema, or other widespread conditions, specialised opinion should be sought in Advice &amp; Guidance attaching clinical photographs.</p>
Viral warts	<p>Children found to have ano-genital warts should be referred to the York 'Child Sexual Assault Assessment Centre' for confirmation of diagnosis.</p> <p>Treatment for Viral Warts is restricted to the minimum eligibility criteria below. This is because most plantar warts can be managed with over the counter topical treatments or by treatments prescribed in Primary Care. Treatment for Viral Warts that do not meet the criteria below are deemed to be cosmetic and will not be funded. Referral to secondary care dermatology should only be made:</p> <ul style="list-style-type: none"> <li>• for ano-genital warts in adults that have failed treatment in the Primary Care setting or Genito-Urinary (GUM) Clinic</li> <li>• for viral warts in immunosuppressed patients</li> <li>• if there is doubt about the diagnosis and concern about possible malignancy</li> <li>• Facial viral warts that have not resolved with an appropriate trial of topical treatment.</li> </ul> <p>Where there are exceptional circumstances, referral should be made to the Individual Funding Request Panel. Viral warts on face where there are physical or mental sequelae should be referred to IFR for funding.</p>
Skin tags (including anal skin tags)	<p>Treatment is not routinely commissioned. Where there is diagnostic uncertainty requesting a specialist opinion by sending photos via Advice and Guidance is recommended. . Where exceptional clinical indications exist (e.g. intractable pruritus ani) then referral to the Individual Funding Request Panel is advised.</p>
Cyst of moll	<p>Not routinely commissioned. Where there is diagnostic uncertainty requesting a specialist opinion by sending photos via Advice and Guidance is recommended.</p>
Cyst of Zeis	<p>Not routinely commissioned. Where there is diagnostic uncertainty requesting a specialist opinion by sending photos via Advice and Guidance is recommended.</p>
Pingueculum	<p>Not routinely commissioned. Where there is diagnostic</p>

	uncertainty requesting a specialist opinion by sending photos via Advice and Guidance is recommended.
Eyelid papillomas and skin tags	Not routinely commissioned. Where there is diagnostic uncertainty requesting an ophthalmologist opinion by sending photos via Advice and Guidance is recommended. See <a href="#">oculoplastic eye problems commissioning statement</a> .
Actinic solar keratosis (AK)	<p>Referral to secondary care for Actinic Keratosis is not expected unless primary care treatments have failed, (guidance on primary care treatment is on the Referral Support Site website under Dermatology).</p> <p>Refer to secondary care for:</p> <ul style="list-style-type: none"> <li>• severe AK when there may be a possibility of invasive malignancy: these are thicker and harder and may have an infiltrated base refer to secondary care where there is diagnostic uncertainty.</li> <li>• failure of 2 different treatments</li> <li>• Immuno-compromised patients</li> </ul>
Pigmented Naevi (moles)	Refer if there is clinical suspicion of malignancy or diagnostic uncertainty.
Lipoma	<p>If size less than 3cm, asymptomatic with no diagnostic uncertainty Surgery is NOT routinely funded for cosmetic reasons and concerns about cosmetic appearance should NOT be referred to secondary care unless there are clinically exceptional circumstances with IFR Panel approval. See Guidance <a href="#">here</a>.</p> <p>Referral for diagnostic uncertainty only</p>
<b>Summary of evidence / rationale</b>	<p>Minor surgery should only be carried out when clinically necessary and after weighing up the risks and benefits.</p> <p>The use of NHS resources to manage benign cosmetic lesions is not a current priority and expectations of such should be discouraged.</p> <p>The risks of carrying out minor surgery on skin lesions include damage to nerves, haemorrhage, failure to achieve wound closure, wound infection, wound dehiscence, over granulation, incomplete excision rate, unsatisfactory scar formation and distortion to local anatomy<sup>1</sup></p> <p>A comparison of minor surgery in primary and secondary care carried out in the South of England suggested that the quality of minor surgery carried out in general practice is not quite as high as that carried out in hospital, but patients prefer the convenience of treatment in General Practice. However, there</p>

	may be clear deficiencies in GPs' ability to recognise malignant lesions, and there may be differences in completeness of excision when compared with hospital doctors <sup>2</sup>
<b>Date effective from</b>	1 <sup>st</sup> May 2019
<b>Date published</b>	30 <sup>th</sup> May 2019
<b>Review date</b>	30 <sup>th</sup> April 2020
<b>Approved by</b>	Vale of York Executive Committee and Scarborough and Ryedale Business Committee
<b>Responsible officer</b>	Simon Cox Director of Acute Commissioning

**References:**

1. Primary Care Dermatology Society - Skin Surgery Guidelines 2007  
[http://www.pcds.org.uk/images/downloads/skin\\_surgery\\_guidelines.pdf](http://www.pcds.org.uk/images/downloads/skin_surgery_guidelines.pdf)
2. S George, et al. (2008) A prospective randomised comparison of minor surgery in primary and secondary care. The MiSTIC trial. Health Technology Assessment 2008; Vol. 12: No. 23.  
[http://www.journalslibrary.nihr.ac.uk/\\_data/assets/pdf\\_file/0006/64905/FullReport-hta12230.pdf](http://www.journalslibrary.nihr.ac.uk/_data/assets/pdf_file/0006/64905/FullReport-hta12230.pdf)
3. NHSE/I.(2018) Evidence-Based Interventions: Consultation Document.  
 Available at: <https://www.england.nhs.uk/wp-content/uploads/2018/06/04-b-pb-04-07-2018-ebi-consultation-document.pdf>. Accessed:18/12/2018