Management of benzodiazepine misuse

Clinical Institute Withdrawal Assessment Scale - benzodiazepines

Client report:

Do you feel	0 Not at all	1	2	3	4 Very much
irritable?		_	_		so
Do you feel	0 Not at all	1	2	3	4 Unable to
fatigued?					function
Do you feel	0 Not at all	1	2	3	4 Very much
tense?					so
Do you have	0 Not at all	1	2	3	4 Unable to
difficulties					concentrate
concentrating?					
Do you have any	0 Not at all	1	2	3	4 No
loss of appetite?					appetite
					unable to
					eat
Do you have any	0 No	1	2	3	4 Intense
numbness or	numbness				burning/
burning on your					numbness
face, hands or					
feet					
Do you feel your	0 No	1	2	3	4 Constant
heart racing?	disturbance				racing
(palpitation)					
Does your head	0 Not at all	1	2	3	4 Severe
feel full or achy?					headache
Do you feel	0 Not at all	1	2	3	4 Severe
muscle aches or					stiffness or
stiffness?					pain
Do you feel	0 Not at all	1	2	3	4 Very much
anxious, nervous					so
or jittery?					
Do you feel	0	1	2	3	4Very much
upset?	Not at all				so
How restful was	0 Very	1	2	3	4 Not al all
your sleep last	restful				
night?					
Do you feel	0 Not at all	1	2	3	4 Very much
weak?					so
Do you think you	0 Very	1	2	3	4 Not at all

didn't have enough sleep last night?	much so				
Do you have any visual disturbances? (sensitivity to light, blurred vision)	0 Not at all	1	2	3	4 Very sensitive to light, blurred vision
Are you fearful?	0 Not at all	1	2	3	4 Very much so
Have you been worrying about possible misfortunes lately?	0 Not at all	1	2	3	4 Very much so

Clinician observations:

Observe behaviour for restlessness and agitation	0 none, normal activity	1	2 restless	3	4 paces back and forth, unable to sit still
Observe tremor	0 No tremor	1 Not visible, can be felt in fingers	2 visible but mild	3 Moderate with arms extended	4 severe, with arms not extended
Observe for sweating – feel palms	0 no sweating visible	1 barely perceptible sweating, palms moist	2 palms and forehead moist, reports armpit sweating		

Interpretation of scores:

1–20 = mild withdrawal

21–40 =moderate withdrawal

41–60 = severe withdrawal

61–80 = very severe withdrawal

Ref: Busto, U.E., Sykora, K. & Sellers, E.M. (1989). A clinical scale to assess benzodiazepine withdrawal. Journal of Clinical Psychopharmacology, 9 (6), 412–416.