

## Referral Support Service

### Breast

#### BO2 Breast Lumps

[Click here for 2WW referral form](#)

##### Definition

Any new lump in the breast

##### Exclude Red Flag Symptoms

- Discrete lump in woman aged 30 or over.
- Associated ulceration, skin nodule, skin distortion or palpable axillary lymph nodes.
- Unilateral firm subareolar mass in men aged 50 or more.
- Associated erythema or redness of the breast tissue.
- Increase in size or change in character of a pre existing or previously investigated lump
- All require urgent referral under Two Week Rule

##### Management

- Asymmetric nodularity should be reassessed after 6 weeks and referred if persistent.
- Young women <35 with long standing painful lumpy breasts and older women with symmetrical nodularity if not a localized abnormality can be managed in Primary Care with reassurance and review.
- Smoking cessation to be encouraged
- Weight loss (if appropriate) to be encouraged

##### Referral Information

###### Information to include in referral letter

- Site and duration of lump
- Any family history of breast or ovarian cancer
- Relevant past medical / surgical history
- Current regular medication
- BMI/ Smoking status

##### Patient Information Leaflets/ PDAs

[www.patient.co.uk/health/breast-lump](http://www.patient.co.uk/health/breast-lump)

Responsible GP: Dr Lorraine Boyd  
Responsible Consultant: Ben Mancy-Jones

Clinical Research & Effectiveness approved: April 2016  
Date published: May 2016  
Next review: May 2018

© NHS Vale of York Clinical Commissioning Group – Version 2

The on-line version is the only version that is maintained. Any printed copies should, therefore, be viewed as 'uncontrolled' and as such may not necessarily contain the latest updates and amendments.

## References

- NICE Referral Advice Updated 22 08 12
- CRUK Guidelines for Referral of Patients with Breast Problems
- SIGN Guideline 84: Management of Breast Cancer in Women